

Status and Plans for the NIMROD Code Development Project

C. R. Sovinec, A. H. Glasser, and R. A. Nebel
Los Alamos National Laboratory

T. A. Gianakon
DRFC/SPPF, CE Cadarache

A. E. Koniges
Lawrence Livermore National Laboratory

M. W. Phillips
Northrup Grumman Corporation

S. J. Plimpton
Sandia National Laboratory

D. D. Schnack
Science Applications International Corporation

and the NIMROD Team

presented at:

1997 International Sherwood Fusion Theory Conference

April 28-30, 1997

Madison, Wisconsin

OUTLINE

- I. NIMROD introduction
 - A. Participants
 - B. Objectives
 - C. Product

- II. Physics Kernel
 - A. Revised formulation
 - B. Development status
 - C. Linear solver

- III. Benchmarking
 - A. Plan
 - B. Linear results for comparison

- IV. Graphical User Interface

- V. Preprocessing

- VI. Postprocessing
 - A. XDRAW
 - B. IBM Data Explorer

OBJECTIVES

1. Model low-frequency phenomena in high-performance tokamaks.
2. Provide a flexible computational tool for alternates.
3. Explore the use of *concurrent engineering* and *quality function deployment* in large-scale code development.

PRODUCT

Non-Ideal MHD with Rotation: Open Discussion

- Time-dependent, time-split, implicit fluid simulation code.
- Realistic, flexible geometry via finite element discretization of the poloidal plane.
- Pseudospectral representation of the periodic direction.
- Block decomposition of the computational domain for parallel computing on message passing architectures.
- Graphical User Interface for easy use.
- Multi-level graphics for routine and publication quality output.

PHYSICS KERNEL

Original numerical formulation:

\mathbf{J}_S equations \rightarrow Ampere's law \rightarrow implicit \mathbf{E} equation.

Unacceptable coupling of shear and compressional Alfvén waves.

Revised formulation:

Generalized Ohm's law \rightarrow Faraday's law \rightarrow implicit \mathbf{B} equation.

Better numerical properties (see Glasser, et al.)

Time-discrete Equations (Cold Plasma)

$$Z_e = -q_i/q_e$$

$$\nu = Z_e m_e/m_i$$

$$m = m_i/Z_e + m_e$$

$$n = n_e, \text{ electron number density}$$

$$\rho = mn = m_i n(1+\nu)/Z_e, \text{ mass density}$$

$$\mathbf{M} = m_i \mathbf{J}_i/q_i + m_e \mathbf{J}_e/q_e, \text{ momentum density}$$

$$f_\Omega = \text{numerical time-centering parameter}$$

$$\frac{\Delta \mathbf{J}_s}{\Delta t} + \frac{f_\Omega q_s}{m_s} \mathbf{B}^* \times \mathbf{J}_s^{n+1} = \frac{n_s q_s^2}{m_s} \mathbf{E} + \frac{(f_\Omega - 1) q_s}{m_s} \mathbf{B}^* \times \mathbf{J}_s^n$$

$$\mathbf{E} = \frac{1}{\epsilon_0 \omega_p^2} \frac{\Delta \mathbf{J}}{\Delta t} + f_\Omega \mathbf{B}^* \times \left(\frac{1}{\rho} \mathbf{M}^{n+1} - \frac{(1-\nu)}{(1+\nu)} \frac{1}{en} \mathbf{J}^{n+1} \right) \\ + (1-f_\Omega) \mathbf{B}^* \times \left(\frac{1}{\rho} \mathbf{M}^n - \frac{(1-\nu)}{(1+\nu)} \frac{1}{en} \mathbf{J}^n \right)$$

$$\mathbf{M}^{n+1} = \mathbf{M}^n + \Delta t (f_\Omega \Delta \mathbf{J} + \mathbf{J}^n) \times \mathbf{B}^*$$

$$\mathbf{E}_{imp} = \frac{1}{\epsilon_0 \omega_p^2} \frac{\Delta \mathbf{J}}{\Delta t} + \frac{f_\Omega^2 \Delta t}{\rho} \left(\mathbf{B}^{*2} - \mathbf{B}^* \mathbf{B}^* \right) \cdot \Delta \mathbf{J} - \frac{(1-\nu)}{(1+\nu)} \frac{f_\Omega}{en} \mathbf{B}^* \times \Delta \mathbf{J}$$

$$\mathbf{E}_{exp} = \frac{1}{\rho} \left[\mathbf{B}^* \times \mathbf{M}^n + f_\Omega \Delta t \left(\mathbf{B}^{*2} - \mathbf{B}^* \mathbf{B}^* \right) \cdot \mathbf{J}^n \right] - \frac{(1-\nu)}{(1+\nu)} \frac{1}{en} \mathbf{B}^* \times \mathbf{J}^n$$

$$\Delta \mathbf{B} + \nabla \times \mathbf{Z} \cdot \nabla \times \Delta \mathbf{B} = -\Delta t \nabla \times \mathbf{E}_{exp}$$

$$\mathbf{Z} = \frac{c^2}{\omega_p^2} \mathbf{I} + \frac{f_\Omega^2 \Delta t^2}{\mu_0 \rho} \left(\mathbf{B}^{*2} - \mathbf{B}^* \mathbf{B}^* \right) - \frac{(1-\nu)}{(1+\nu)} \frac{f_\Omega \Delta t}{\mu_0 en} \mathbf{B}^* \times \mathbf{I}$$


Present Status

- Revised formulation including conforming/nonconforming approximation options.
- Two types of grid blocks: 1) rectangular blocks of structured quadrilateral cells, 2) blocks of unstructured triangles.
- Cold plasma equations (except semi-implicit term for Hall dynamics).
- Nonzero electrical resistivity (implicit).
- F90 conjugate gradient solver with diagonal preconditioning for blocks of triangles and either diagonal or direct, domain-decomposition preconditioning on rectangular blocks.
- Message passing for parallel computing (see Plimpton, et al.).

In-progress and Planned Development

- Multiple toroidal modes and FFTs for pseudospectral operations.
- Pressure evolution equation including anisotropic thermal conduction.
- Explicit advection terms.
- Neoclassical closure terms.
- Arbitrary unit vectors.
- $\nabla \cdot \mathbf{B}$ cleaning.
- Multi-grid preconditioning for cg solver.

NIMROD Physics Kernel Development



Force-free, Ideal Linear Results for Comparison
(Computed by S. A. Galkin, Keldysh Institute of Applied
Mathematics with the TWIST-R and RESTOR codes.)

BENCHMARKING

NIMROD VALIDATION PLAN

(DRAFT 9/20/96 Rev. 1)

I. Stable linear oscillations (MHD limit only)

A. Uniform B_{ϕ} , no current

1. Zero beta, 2-D

a. Compressional Alfvén waves, plasma oscillations, whistler waves, all cases

have perfectly conducting plasma

i) Rectangular region - frequency known

analytically * Uniform grid

- Explicit

- Implicit

* Non uniform grid

- Explicit

- Implicit

ii) Cylindrical region

* Multiple blocks with seams

2. Zero beta, 3-D

a. Compressional Alfvén waves, plasma oscillations, whistler waves - $n=0$ -

Repeat the above test to assure there are no surprises.

b. Torsional Alfvén waves, $n > 0$ - Compare frequency with analytic (?) solutions and

with GATO. (Initialize with GATO

eigenfunction?) Both explicit and implicit cases. i) Single block grid
ii) Multiple block grid with seams

3. Finite beta, 2-D

- a. Electron sound waves (turn off ions)
 - b. Ion sound waves (both species on)
 - c. Drift waves
4. Finite beta, 3-D redo 2-D cases

B. Force-free toroidal equilibria - Comparison with GATO

frequencies and eigenfunctions. (Are there any analytic equilibria of this sort?)

II. Linear unstable modes, 3D (Ideal MHD limit)

A. Internal modes (perfectly conducting boundary at separatrix).

- 1. Force-free toroidal equilibria (are there any that we understand?)
- 2. Finite beta

a. Solov'ev equilibrium - Internal kink modes

- i) Berger, et al, paper (growth rates and eigenfunctions)

- ii) comparison of marginal points with Berger, et al., and DCON

- b. Numerical equilibria - Compare growth rates and eigenfunctions with GATO, Keldish codes, marginal points with DCON

- i) DIIID

- ii) TFTR

- iii) ITER
- iv) etc, etc, etc.....

B. External modes - vacuum region - Comparisons

with GATO.

III. Linear unstable modes (Resistive MHD)

- A. Compare with Carreras, et al ($n = 2$ tearing mode in a circular cross section torus) - Growth rate and eigenfunction
 - B. Any other known toroidal solutions (Keldish codes?)
 - C. Marginal points (DCON?)
- IV. Linear unstable modes with 2-Fluid effects
- A. Cylindrical case (Pic3d comparison)
 1. Ideal (1,1) mode
 2. Resistive (2,1) mode

B. Toroidal case (QIP3D

comparison)

1. Resistive (1,1)

mode

2. Resistive (2,1)

mode

V. Nonlinear evolution (Ideal and resistive MHD) - We need to identify a D nonlinear codes.

A. Internal modes???

B. External modes (requires moving separatrix)???

VI. Thermal Conductivity - Test pollution of perpendicular transport in highly anisotropic cases. Impose islands and see how well temperature conforms to the flux surfaces.

VII. Nonlinear evolution with 2-Fluid effects

A. Cylindrical case (Pic3d

comparison)

1. Ideal (1,1) mode

2. Resistive (2,1) mode

B. Toroidal case (QIP3D

comparison)

1. Resistive (1,1) mode

2. Resistive (2,1) mode

VIII. Neo-Classical effects - We need to define a suite of test problems here, too.

IX. Comparison with experiment - We need help here, too.

Shear Alfven Wave Tests

- Uniform rectangular grid, doubly periodic, 1m on a side.
- Neither \mathbf{k} nor \mathbf{B} are aligned with the grid ($k_x=k_y=2\pi$, $k_z=2^{3/2}\pi$).

formulation	conform	k-par/k	grid	v_{phase}/v_A
E	y	10%	9x9	1.67
E	y	10%	17x17	1.22
E	n	10%	9x9	1.25
E	n	10%	17x17	1.07
E	y	5%	9x9	2.81
E	y	5%	17x17	1.69
E	n	5%	9x9	1.47
E	n	5%	17x17	1.14
E	y	2.5%	9x9	5.35
E	y	2.5%	17x17	2.91
E	n	2.5%	9x9	1.89
E	n	2.5%	17x17	1.25
B	y	10%	8x8	0.984
B	y	10%	16x16	1.00
B	n	10%	8x8	1.10
B	n	10%	16x16	1.02
B	y	5%	8x8	0.974
B	y	5%	16x16	1.00
B	n	5%	8x8	1.36
B	n	5%	16x16	1.09
B	y	2.5%	8x8	0.95
B	y	2.5%	16x16	1.00
B	n	2.5%	8x8	1.83
B	n	2.5%	16x16	1.22

GRAPHICAL USER INTERFACE

(NIMEXEC and supporting scripts)

Tcl scripts presently provide:

- job status information and control
- convenient input file editing
- links to preprocessing and postprocessing codes

Future developments will allow remote job control.

See demonstration.

PREPROCESSING

(NIMSET & FLUXGRID)

- Read equilibrium code output.
- Generate grid based on flux surfaces.
- Generate block connections and vertex connections within blocks of unstructured triangles.
- Initialize physical fields such as \mathbf{B}_0 and momentum density perturbations.
- Write an initial dump file.

POSTPROCESSING

Routine graphics (XDRAW):

- Grid plots
- Spatial slicing
- Slices showing temporal dependence
- Time-history traces
- Contour plotting of multi-block problems is being developed by N. Popova

IBM Data Explorer

- Multi-dimensional movies
- Sophisticated, 3D data rendering available

See demonstration.